

2579 E17th St Suite 11, Brooklyn, NY 11235 (718) 382-3444 www.LuxDen.com

Perio Surgery - Aftercare

Periodontal surgery

You were given an anesthetic to ensure your comfort. This anesthetic typically leaves your lips, teeth and tongue feeling numb after the appointment. For this reason, you should avoid any chewing for two hours following surgery, or until the numbness has worn off completely.

- > Apply an ice pack 20 min off, 20 min off for six hours following surgery to decrease pain and swelling.
- > Keep your fingers and tongue away from the parts of your gums that were affected by the surgery.
- > Do not rinse your mouth the day of the surgery. Twenty-four hours after surgery, you can rinse gently with mouthwash or a warm salt water solution. Dissolve one teaspoon of salt in one cup of warm water, and gently swish, then carefully spit. Three times per day is sufficient.
- > It's normal to experience some discomfort for a few days following surgery. To avoid nausea, don't take pain medication on an empty stomach.
- > Relax as much as possible and avoid all strenuous activities for the first 24 hours following surgery.
- > Once the numbness has worn off, you should eat. Nourishment is important to the healing process. Limit your diet to soft foods like yogurt, soft soup, ice cream, or soft cooked eggs for the first 48 hours.
- * Drink at least eight large glasses of water or diluted fruit juice each day.

- > Keep your head elevated with pillows to control the bleeding. We will give you a supply of gauze sponges to place over the bleeding area. Change them as necessary, and use them until the bleeding stops completely. You can also bite gently but firmly on a moist tea bag for 20 minutes if bleeding continues.
- > Continue cleaning the other areas of your MOUth, flossing daily and brushing at least twice a day as you would normally.

Call our office is bleeding persists or "CineaS 2S•



Avoid Spicy Food



Temporary Soft Diet



Take All Medications